



# Framework and guidelines for implementing the proposed IUCN Environmental Impact Classification for Alien Taxa (EICAT)

Charlotte L. Hawkins<sup>1</sup>, Sven Bacher<sup>2</sup>, Franz Essl<sup>3</sup>, Philip E. Hulme<sup>4</sup>, Jonathan M. Jeschke<sup>5,6</sup>, Ingolf Kühn<sup>7,8</sup>, Sabrina Kumschick<sup>9,10</sup>, Wolfgang Nentwig<sup>11</sup>, Jan Pergl<sup>12</sup>, Petr Pyšek<sup>12,13</sup>, Wolfgang Rabitsch<sup>14</sup>, David M. Richardson<sup>9</sup>, Montserrat Vilà<sup>15</sup>, John R. U. Wilson<sup>9,10</sup>, Piero Genovesi<sup>16</sup> and Tim M. Blackburn<sup>1,17,18,\*</sup>

<sup>1</sup>Department of Genetics, Evolution & Environment, Centre for Biodiversity & Environment Research, Darwin Building, UCL Gower Street, London WC1E 6BT, UK,

<sup>2</sup>Department of Biology, Unit Ecology & Evolution, University of Fribourg, Chemin du Musée 10, Fribourg 1700, Switzerland,

<sup>3</sup>Division of Conservation Biology, Vegetation and Landscape Ecology, University of Vienna, Rennweg 14, Vienna 1030, Austria, <sup>4</sup>The Bio-Protection Research Centre, Lincoln University, PO Box 84, Christchurch New Zealand, <sup>5</sup>Leibniz-

Institute of Freshwater Ecology and Inland Fisheries (IGB), Müggelseedamm 310, Berlin 12587, Germany, <sup>6</sup>Department of Biology, Chemistry, Pharmacy, Institute of Biology, Freie Universität Berlin, Königin-Luise-Str.

1-3, Berlin 14195, Germany, <sup>7</sup>Department of Community Ecology, UFZ, Helmholtz Centre for Environmental Research, Theodor-Lieser-Str. 4, Halle 06120, Germany, <sup>8</sup>German

Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, Leipzig 04103, Germany, <sup>9</sup>Department of Botany and Zoology, Centre for Invasion

Biology, Stellenbosch University, Private Bag X1, Matieland 7602, South Africa, <sup>10</sup>Invasive Species Programme, South African National Biodiversity Institute, Kirstenbosch National

Botanical Gardens, Claremont 7735, South Africa, <sup>11</sup>Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, Bern

CH-3012, Switzerland, <sup>12</sup>Department of Invasion Ecology, Institute of Botany, The Czech Academy of Sciences, Průhonice CZ-252 43, Czech Republic, <sup>13</sup>Department of Ecology, Faculty of Science, Charles

## ABSTRACT

Recently, Blackburn *et al.* (2014) developed a simple, objective and transparent method for classifying alien taxa in terms of the magnitude of their detrimental environmental impacts in recipient areas. Here, we present a comprehensive framework and guidelines for implementing this method, which we term the Environmental Impact Classification for Alien Taxa, or EICAT. We detail criteria for applying the EICAT scheme in a consistent and comparable fashion, prescribe the supporting information that should be supplied along with classifications, and describe the process for implementing the method. This comment aims to draw the attention of interested parties to the framework and guidelines, and to present them in their entirety in a location where they are freely accessible to any potential users.

## Keywords

alien, environment, impact, IUCN.

University in Prague, Viničná 7, CZ-128 44

Praha 2, Prague Czech Republic,

<sup>14</sup>Department of Biodiversity and Nature  
Conservation, Environment Agency Austria,  
Spittelauer Lände 5, Vienna 1090, Austria,

<sup>15</sup>Estación Biológica de Doñana (EBD-  
CSIC), Avda. Américo Vespucio s/n Isla de  
la Cartuja, Sevilla 41092, Spain, <sup>16</sup>ISPRA,  
Institute for Environmental Protection and  
Research and Chair IUCN SSC Invasive  
Species Specialist Group, Via Vitaliano  
Brancati 48, Rome 00144, Italy,

<sup>17</sup>Distinguished Scientist Fellowship Program,  
King Saud University, Riyadh 1145, Saudi  
Arabia, <sup>18</sup>Environment Institute, School of  
Earth & Environmental Sciences, University  
of Adelaide, Adelaide SA 5005, Australia

\*Correspondence: Tim M. Blackburn,  
Department of Genetics, Evolution &  
Environment, Centre for Biodiversity &  
Environment Research, Darwin Building,  
UCL, Gower Street, London WC1E 6BT, UK.  
E-mail: t.blackburn@ucl.ac.uk

Many invasive alien taxa have substantial effects on the ecosystems into which they have been introduced, including significant changes in native species extinction probabilities, genetic composition, behaviour patterns, richness and abundance, as well as altering phylogenetic and taxonomic diversity, trophic networks, ecosystem productivity, nutrient cycling, geomorphology, hydrology, habitat structure and various components of disturbance regimes (Brooks *et al.*, 2004; Hendrix *et al.*, 2008; Suarez & Tsutsui, 2008; Kenis *et al.*, 2009; Winter *et al.*, 2009; Vilà *et al.*, 2011; Pyšek *et al.*, 2012; Ricciardi *et al.*, 2013; Fei *et al.*, 2014). As a result, most scientists and conservation organizations consider invasive alien species to be undesirable (Lambertini *et al.*, 2011), and considerable resources have been devoted towards preventing or mitigating their impacts. Nevertheless, impacts vary greatly among alien taxa and among ecosystems, habitats or native species in the recipient geographic range; many notable impacts only become obvious or significantly influential long after the onset of invasion (Williamson & Fitter, 1996; Graves & Shapiro, 2003; Sogge *et al.*, 2008; Stromberg *et al.*, 2009; Chiba, 2010; Schlaepfer *et al.*, 2011; Pyšek *et al.*, 2012; Strayer, 2012). Thus, it has long been recognized that there is a critical need for the capacity to evaluate, compare and predict the magnitudes of the impacts of different alien taxa, in order to determine and prioritize appropriate actions.

In response to this need, Blackburn *et al.* (2014) recently developed a simple, objective and transparent method for classifying alien taxa in terms of the magnitude of their detrimental environmental impacts in recipient areas. We refer to this scheme as the Environmental Impact Classifica-

tion for Alien Taxa, or EICAT. Alien taxa are classified into one of five 'impact' categories depending on the level of biological organization (individual, population or community) impacted, with the mechanisms by which the impacts occur aligned with those identified in the International Union for Conservation of Nature (IUCN) Global Invasive Species Database (<http://www.issg.org/database/welcome>). If applied consistently, EICAT will help to (1) identify those taxa that have different levels of environmental impact, distinguishing taxa causing impacts of low concern from invasive alien taxa with significant deleterious effects (those with negative impacts, *sensu* the definition from the Convention on Biological Diversity); (2) facilitate comparisons of the level of impact by alien taxa among regions and taxonomic groups; (3) facilitate predictions of potential future impacts of alien taxa in the target region and elsewhere; (4) aid in prioritization of management actions; and (5) aid in evaluation of management methods. It is envisaged that EICAT will be used by scientists, land managers and conservation practitioners as a tool to gain a better understanding of the magnitude of impacts caused by different alien taxa and to inform the prioritization, implementation and evaluation of management policies and actions within existing international agreements and statutes. In this regard, one aim is to have the scheme adopted as the official mechanism by which IUCN classifies the deleterious environmental impacts of those alien taxa that have become globally or regionally invasive.

A key requirement for any standardized scheme adopted under the auspices of IUCN (or any other major administrative organization overseeing international conventions and policy implementation) is the existence of a clear framework

and guidelines for implementing the scheme. To this end, we have drafted just such a framework and guidelines for implementing the EICAT scheme: the primary aim of this contribution is to draw the attention of interested parties to the framework and guidelines, and to present them in their entirety in a location where they are freely accessible to any potential users. Hence, we have appended them to this comment as supplementary online information (see Appendix S1 in Supporting Information).

To ensure that the maximum utility can be gained from any classifications carried out using EICAT, it is important that studies apply the scheme in a consistent and comparable fashion, and that consistent and comprehensive supporting information is supplied along with classifications. Therefore, the framework (1) defines some of the key concepts required to use EICAT; (2) describes the categories and criteria for classifying alien taxa, the guidelines for transferring taxa between categories and the taxonomic and geographic scope of the scheme; (3) provides a method for dealing with uncertainty over the correct classification, and in particular uncertainty that arises from the need to interpret evidence generated at spatial scales that are often very different to the spatial scales over which native communities can be characterized; (4) prescribes essential and recommended documentation that should or could be provided in support of classifications; and (5) describes the process for implementing these guidelines.

We have intended the EICAT framework (see Appendix S1) to be comprehensive, and so it reproduces the key elements of the classification scheme presented in Blackburn *et al.* (2014). However, it describes two changes to the scheme as originally published that are important to note. First, the EICAT framework describes parallel classification systems to capture both the maximum impact ever recorded for an alien taxon at a point in time, and the current impact level caused by that taxon at the same location or elsewhere in the geographical area of interest (e.g. globally, regionally, nationally). This addition ensures that EICAT captures the maximum recorded impact of an alien taxon introduced somewhere in the geographical area of interest, which may be especially important where the current impact is not as high as in the past, perhaps due to mitigation measures or spontaneous changes (Dostál *et al.*, 2013). Instigating parallel classifications for current and the maximum impacts ever recorded also helps in indicating any changes in current environmental impact through successive assessments (i.e. as a result of the invasion proceeding, or mitigation measures taking effect).

Second, the framework alters the two-letter abbreviation codes for the five categories of impact, to try to reduce the possibility for ambiguity in the original designations (and so that the codes increase in alphabetical order). These now become Minimal Concern (MC), Minor (MN), Moderate (MO), Major (MR) and Massive (MV).

The process for implementing the EICAT identifies an EICAT Unit, consisting of members of the IUCN Invasive

Species Specialist Group, whose function it is to oversee the entire process, and to check each assessment to ensure consistency. The Unit is also intended to coordinate the reporting of status and trends in impacts as documented by the EICAT process and oversee any proposals for changes or revisions to the framework and guidelines. Impact classification assessments should be independently reviewed by experts, to check whether the assessment has been carried out appropriately as described in the framework and guidelines, before being submitted to the EICAT Unit for final ratification. The aim is to have accepted assessments published on the Global Invasive Species Database, where they will be freely available to the community of relevant stakeholders. Anyone who has carried out environmental impact classifications under the EICAT scheme should contact one of the current members of the EICAT Unit (see Appendix S1, section 7), to discuss the formal adoption of their assessment(s) under this scheme.

## ACKNOWLEDGEMENTS

This study is an output of the working group sImpact that formed at a workshop at sDiv, the Synthesis Centre within the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (DFG; through grant no. FZT 118). Financial support was also provided by the King Saud University Distinguished Scientist Fellowship Program (to CLH and TMB); sDiv and GEO BON; the ERA-Net BiodivERsA (project FFII, with the national funder DFG, grant JE 288/7-1) and the DFG (grant JE 288/9-1) (to JMJ); the Swiss National Science Foundation and the Drakenstein Trust through the DST-NRF Centre of Excellence for Invasion Biology, and The South African Department of Environmental Affairs through its funding of SANBI's Invasive Species Programme (to SK); long-term research development project RVO 67985939 and Praemium Academiae award from The Czech Academy of Sciences (to PP); the National Research Foundation, South Africa (grants 85417 to DMR and 86894 to JRW); and the Severo Ochoa Program for Centres of Excellence in R+D+I (SEV-2012-0262) from the Spanish Ministry of Economy and Competitiveness (to MV). Support from the EU COST Action TD1209 Alien Challenge is also gratefully acknowledged. We thank Myriam Du Mortier, Craig Hilton-Taylor, Donald Hobern, Nick Holmes, Melodie McGeoch, Sean Murphy, Jon-Paul Rodriguez, Riccardo Scaler, Junko Shimura and Marten Winter for comments and discussion that substantially improved the framework and guidelines document.

## REFERENCES

- Blackburn, T.M., Essl, F., Evans, T., Hulme, P.E., Jeschke, J. M., Kühn, I., Kumschick, S., Marková, Z., Mrugała, A., Nentwig, W., Pergl, J., Pyšek, P., Rabitsch, W., Ricciardi,

- A., Richardson, D.M., Sendek, A., Vilá, M., Wilson, J.R.U., Winter, M., Genovesi, P. & Bacher, S. (2014) A unified classification of alien species based on the magnitude of their environmental impacts. *PLoS Biology*, **12**, e1001850.
- Brooks, M.L., D'Antonio, C.M., Richardson, D.M., Grace, J.B., Keeley, J.E., DiTomaso, J.M., Hobbs, R.J., Pellant, M. & Pyke, D. (2004) Effects of invasive alien plants on fire regimes. *BioScience*, **54**, 677–688.
- Chiba, S. (2010) Invasive non-native species' provision of refugia for endangered native species. *Conservation Biology*, **24**, 1141–1147.
- Dostál, P., Müllerová, J., Pyšek, P., Pergl, J. & Klinerová, T. (2013) The impact of an invasive plant changes over time. *Ecology Letters*, **16**, 1277–1284.
- Fei, S., Phillips, J. & Shouse, M. (2014) Biogeomorphic impacts of invasive species. *Annual Review of Ecology, Evolution, and Systematics*, **45**, 69–87.
- Graves, S.D. & Shapiro, A.M. (2003) Exotics as host plants of the California butterfly fauna. *Biological Conservation*, **110**, 413–433.
- Hendrix, P.F., Callahan, M.A., Drake, J.M., Huang, C.-Y., James, S.W., Snyder, B.A. & Zhang, W. (2008) Pandora's Box contained bait: the global problem of introduced earthworms. *Annual Review of Ecology, Evolution, and Systematics*, **39**, 593–613.
- Kenis, M., Auger-Rozenberg, M.-A., Roques, A., Timms, L., Péré, C., Cock, M.W., Settele, J., Augustin, S. & Lopez-Vaamonde, C. (2009) Ecological effects of invasive alien insects. *Ecological impacts of non-native invertebrates and fungi on terrestrial ecosystems SE - 3* (ed. by D. Langor and J. Sweeney), pp. 21–45. Springer, Netherlands.
- Lambertini, M., Leape, J., Marton-Lefevre, J., Mittermeier, R., Rose, M., Robinson, J., Stuart, S., Waldman, B. & Genovesi, P. (2011) Invasives: a major conservation threat. *Science*, **333**, 404–405.
- Pyšek, P., Jarošík, V., Hulme, P.E., Pergl, J., Hejda, M., Schaffner, U. & Vilá, M. (2012) A global assessment of invasive plant impacts on resident species, communities and ecosystems: the interaction of impact measures, invading species' traits and environment. *Global Change Biology*, **18**, 1725–1737.
- Ricciardi, A., Hoopes, M.F., Marchetti, M.P. & Lockwood, J.L. (2013) Progress towards understanding the ecological impacts of nonnative species. *Ecological Monographs*, **83**, 263–282.
- Schlaepfer, M.A., Sax, D. & Olden, J.D. (2011) The potential conservation value of non-native species. *Conservation Biology*, **25**, 428–437.
- Sogge, M.K., Sferri, S.J. & Paxton, E.H. (2008) Tamarix as habitat for birds: implications for riparian restoration in the southwestern United States. *Restoration Ecology*, **16**, 146–154.
- Strayer, D.L. (2012) Eight questions about invasions and ecosystem functioning. *Ecology Letters*, **15**, 1199–1210.
- Stromberg, J.C., Chew, M.K., Nagler, P.L. & Glenn, E.P. (2009) Changing perceptions of change: the role of scientists in Tamarix and river management. *Restoration Ecology*, **17**, 177–186.
- Suarez, A.V. & Tsutsui, N.D. (2008) The evolutionary consequences of biological invasions. *Molecular Ecology*, **17**, 351–360.
- Vilá, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J.L., Pergl, J., Schaffner, U., Sun, Y. & Pyšek, P. (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecology Letters*, **14**, 702–708.
- Williamson, M. & Fitter, A. (1996) The varying success of invaders. *Ecology*, **77**, 1661–1666.
- Winter, M., Schweiger, O., Klotz, S., Nentwig, W., Andriopoulos, P., Arianoutsou, M., Basnou, C., Delipetrou, P., Didziulis, V., Hejda, M., Hulme, P.E., Lambdon, P.W., Pergl, J., Pyšek, P., Roy, D.B. & Kühn, I. (2009) Plant extinctions and introductions lead to phylogenetic and taxonomic homogenization of the European flora. *Proceedings of the National Academy of Sciences USA*, **106**, 21721–21725.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

**Appendix S1.** The complete version of the framework and guidelines for implementing the Environmental Impact Classification for Alien Taxa (EICAT).

## BIOSKETCH

**Tim M. Blackburn** is an ecologist with interests in human-mediated biological invasions, extinction and other topics relating to the large-scale distribution and abundance of species. He mainly works on birds.

Editor: Richard Duncan